

Rolf Rose,^{a‡} Franziska
Huttenlocher,^{b‡} Anna Cedzich,^b
Markus Kaiser,^a Andreas
Schaller^{b*} and Christian
Ottmann^{a*}

^aChemical Genomics Centre, Otto-Hahn-Strasse
15, 44227 Dortmund, Germany, and ^bInstitute
of Plant Physiology and Biotechnology,
University of Hohenheim, Emil-Wolff-Strasse
25, 70593 Stuttgart, Germany

‡ These authors contributed equally.

Correspondence e-mail:
schaller@uni-hohenheim.de,
christian.ottmann@cgc.mpg.de

Received 17 March 2009
Accepted 14 April 2009

Purification, crystallization and preliminary X-ray diffraction analysis of a plant subtilase

The subtilase SBT3 from *Solanum lycopersicum* (tomato) was purified from a tomato cell culture and crystallized using the sitting-drop vapour-diffusion method. A native data set was collected to 2.5 Å resolution at 100 K using synchrotron radiation. For experimental phasing, CsCl-derivative and tetrakis-(acetoxymethyl)mercuri-methane (TAMM) derivative crystals were employed for MIRAS phasing. Three caesium sites and one TAMM site were identified, which allowed solution of the structure.

1. Introduction

Subtilases belong to the S8 family of serine proteases (<http://merops.sanger.ac.uk>), which are found in archaea, bacteria, fungi and yeasts as well as in lower and higher eukaryotes. Their active sites display a catalytic triad comprised of aspartate, histidine and serine residues. Subtilases have been grouped into six different families: subtilisins, thermitases, proteinases K, lantibiotic peptidases, kexins and pyrolysins (Siezen & Leunissen, 1997). Only three of them, the kexin, pyrolysin and proteinase K families, are represented in eukaryotes. Kexin, the first eukaryotic subtilase to be identified, processes the precursors of α -mating factor and killer toxin in yeast (Fuller *et al.*, 1989). Nine subtilases have since been identified in mammals, seven of which are kexin-related and are involved in the highly specific processing of precursor proteins. The two remaining subtilases, S1P and PCSK9, belong to the pyrolysin and proteinase K subfamilies, respectively. Many more subtilases can be found in plants compared with animals and most of them have been assigned to the pyrolysin family (Siezen & Leunissen, 1997). The 56 subtilase genes of the model plant *Arabidopsis thaliana*, for example, are divided into six subfamilies, with five being similar to the pyrolysins and one more closely resembling animal kexins (Rautengarten *et al.*, 2005). 63 subtilase genes have been identified in rice (Tripathi & Sowdhamini, 2006) and 15 subtilases have been cloned from tomato (Meichtry *et al.*, 1999). Plant subtilases have been implicated in general protein turnover (Yamagata *et al.*, 1994), in the regulation of plant development (Berger & Altmann, 2000), in biotic and abiotic stress responses (Vera & Conejero, 1988; Liu *et al.*, 2007) and in the processing of mitogenic peptide growth factors (Srivastava *et al.*, 2008).

Despite the prevalence and importance of these proteases in plants, no structural data are available for plant subtilases. We therefore purified and crystallized the tomato subtilase SBT3 (GenBank accession No. AJ006380.1; amino acids 113–761, 69.47 kDa) from a tomato cell culture and solved the structure *via* MIRAS phasing.

2. Methods and results

2.1. Generation of a cell line overexpressing SBT3

A tomato cell-suspension culture was maintained in a Murashige–Skoog-type medium containing 5.0 mg l⁻¹ 1-naphthylacetic acid as described in Felix & Boller (1995). 6 d after subculture, cells were transformed by particle bombardment. Briefly, a 50 ml culture was allowed to sediment and the culture supernatant was replaced with



Table 1

Data-collection statistics.

Values in parentheses are for the outer shell.

	Native	CsCl derivative	TAMM derivative
Wavelength (Å)	0.9796	1.5	1.008
Temperature (K)	100	100	100
Space group	$P4_32_12$	$P4_32_12$	$P4_32_12$
Unit-cell parameters			
a (Å)	143.5	143.9	144.6
b (Å)	143.5	143.9	144.6
c (Å)	196.9	195.1	199.1
α (°)	90	90	90
β (°)	90	90	90
γ (°)	90	90	90
Matthews coefficient (Å ³ Da ⁻¹)	3.63	3.63	3.63
Solvent content (%)	66.13	66.13	66.13
No. of molecules in ASU	2	2	2
Resolution range (Å)	25–2.5 (2.6–2.5)	20–2.6 (3.0–2.6)	15–3.2 (3.4–3.2)
Observed reflections	814751 (91136)	1017071 (433517)	168962 (28272)
Unique reflections	71521 (7807)	135035 (57041)	65272 (10937)
Completeness (%)	100 (100)	99.7 (99.9)	98.8 (99.6)
$R_{\text{merge}}^{\dagger}$	8.3 (32.7)	9.3 (25.9)	7.7 (23.4)
Average $I/\sigma(I)$	17.15 (4.32)	12.13 (5.11)	14.16 (4.78)

$\dagger R_{\text{merge}} = \frac{\sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl) \rangle|}{\sum_{hkl} \sum_i I_i(hkl)}$, where $I_i(hkl)$ is the i th observation of reflection hkl and $\langle I(hkl) \rangle$ is the weighted average intensity for all observations i of reflection hkl .

the same volume of fresh medium supplemented with 10% sucrose. After 30 min of gentle agitation, the cells were sedimented again and 2 ml of packed cell volume was distributed onto agar (0.6%) plates prepared with the same high osmotic strength medium. The cells were then bombarded with spherical gold particles (10 µg; 1.5–3 µm; Aldrich Chemical Co.) coated with 0.6 µg of the linearized (*Sac*I) SBT3 expression construct (*i.e.* the SBT3 cDNA under control of the constitutive CaMV 35S promoter) using a particle inflow gun (Bio-Rad, München). The cells were allowed to recover for 1 d and were then washed from the agar plate into a six-well tissue-culture dish using 6 ml culture medium. After one week at 298 K on an orbital shaker (100 rev min⁻¹), transformed clones were selected on culture plates supplemented with 75 µg ml⁻¹ kanamycin. After 2–3 weeks, when initial callus growth was observed, the calluses were transferred individually to new plates and maintained on the same selection medium. Individual clones were analyzed for stable integration of the transgene by Southern blot analysis and lines with high levels of SBT3 expression were selected by Western blot analysis. A cell-suspension culture was established from the line showing the highest


level of SBT3 expression and maintained as described for the wild-type culture.

2.2. Purification of recombinant SBT3

200 ml culture medium in 1 l Erlenmeyer flasks was inoculated with a 1/10 volume of the SBT3-expressing cell line on the day of subculture. After 8 d at 298 K and 100 rev min⁻¹, the culture supernatant was harvested by vacuum filtration, chilled to 277 K and subjected to fractionated ammonium sulfate precipitation. The precipitate (60–85% saturation) from 6 l culture supernatant was resuspended in 60 ml buffer *A* (25 mM Na₂HPO₄/NaH₂PO₄ pH 7.0, 5 mM EDTA), centrifuged to remove insoluble polysaccharides and cellular debris (2600g, 15 min) and dialysed against two changes of 4 l each of buffer *A* at 277 K. Proteins were then absorbed onto a cation-exchange matrix (SP Sepharose FF, GE Healthcare, Freiburg) equilibrated in buffer *A* using a batch procedure. After washing in buffer *A*, proteins were eluted from the matrix in two consecutive steps using 1 M NaCl in buffer *A*. Eluates were combined and dialysed twice against 4 l buffer *B* (25 mM Tris–HCl pH 9.2, 5 mM EDTA), the dialysate was cleared by filtration (0.45 µm cellulose acetate; Millipore, Schwalbach) and the volume was reduced by ultrafiltration (10 000 molecular-weight cutoff, Vivascience, Hanover). The sample was subjected to anion-exchange chromatography on Resource Q (6 ml column on an ÄKTA purifier chromatography system; GE Healthcare) equilibrated in buffer *B*. The protein was eluted with a linear gradient of NaCl in buffer *B* (0–1 M in 120 ml) at a flow rate of 3 ml min⁻¹. Fractions were assayed for SBT3 activity using an internally quenched fluorogenic peptide substrate [10 µM Abz-SKRDPKMQTD(NO₂)Y; JPT Peptide Technologies, Berlin] in an assay volume of 200 µl 50 mM Tris–HCl pH 8.0 and fluorescence was monitored continuously in a Cary Eclipse spectrofluorimeter (Varian, Darmstadt; $\lambda_{\text{ex}} = 320$ nm, $\lambda_{\text{em}} = 420$ nm). The fractions displaying the highest activity were combined, the volume of the pooled sample was reduced to result in a protein concentration of 10–12 mg ml⁻¹ and the buffer was exchanged to 25 mM Tris–HCl pH 7.0 by ultrafiltration. From 6 l of cell-culture supernatant, the procedure yielded 1.2 mg pure SBT3 as judged by SDS–PAGE.

2.3. Crystallization

In order to crystallize SBT3, initial screenings employing Crystal Screen, Crystal Screen 2 and several Grid Screens from Hampton


Figure 1

(*a*) Crystals of SBT3 grown in 0.1 M Bicine pH 9.0, 10% (w/v) PEG 6000 at 293 K. These crystals did not diffract beyond 6 Å resolution. (*b*) Crystals of SBT3 grown in 0.1 M Bicine pH 8.4, 10% (w/v) PEG 6000, 2.0 M NaCl at 277 K. These crystals diffracted to 4 Å resolution. The scale bar is 100 µm in length.

Research were performed at 293 K by vapour diffusion. The protein concentration in the setups was 11 mg ml⁻¹; 100 nl protein solution was mixed with 100 nl reservoir solution in sitting drops using a Mosquito pipetting robot (TTP LabTech, Melbourn, UK). The sitting drops were equilibrated against reservoirs with a volume of 70 µl. After 2 d, small rosette-like crystals appeared in several conditions. Even the most promising of these fragile crystals [Fig. 1*a*; grown in Grid Screen PEG 6000 condition B6; 0.1 M Bicine pH 9.0, 10% (w/v) PEG 6000] were very thin in one dimension and did not diffract beyond 6 Å resolution. By lowering the pH to 8.4, adding NaCl to a final concentration of 2.0 M and lowering the temperature to 277 K, the crystal morphology and diffraction behaviour could be enhanced significantly (Fig. 1*b*). However, these crystals still did not diffract beyond 4.0 Å resolution. Finally, the addition of 10 mM hexamine-




Figure 2
Crystals of SBT3 grown in 0.1 M Bicine pH 8.4, 10% (w/v) PEG 6000, 2.0 M NaCl and 0.01 M hexamminecobalt(III) chloride at 277 K. These crystals diffracted to 2.5 Å resolution and were used for experimental phasing by fast-soaking them with 0.2 M CsCl or 0.01 M TAMM. The scale bar is 100 µm in length.




Figure 3
A typical X-ray diffraction pattern of an SBT3 crystal fast-soaked in 0.2 M CsCl. The diffraction image was collected on a MAR225 CCD detector. The oscillation range was 0.5° and the crystal-to-detector distance was 270 mm.

Table 2
Phasing statistics.

Phasing took place with two derivatives by MIRAS using *SOLVE* in the resolution range 3.50–20.00 Å. All data sets were collected on SLS beamline PXII and were combined with *CAD* and scaled with *SCALEIT* prior to use in *SOLVE*.

Phasing set	Native	TAMM	CsCl
Radiation source	SLS PXII	SLS PXII	SLS PXII
Preparation	—	24 h soak, 10 mM in cryosolution	20 min soak, 200 mM in cryosolution
Heavy-atom location method	—	INANO, ANOREFINE	INANO, NOANOREFINE
Phasing power (acentric/centric)	—	1.48/1.11	0.73/0.56
Figure of merit (all data)	—	0.29	0.29

Site	Occupancy	x	y	z	B _{iso} (Å ²)
Cs	0.9622	0.5162	0.0548	0.0660	60.0000
Cs	0.3096	0.2603	0.0532	0.0563	34.0941
Cs	0.3968	0.3188	0.0662	0.0474	23.2865
TAMM	0.4994	0.2219	0.0871	0.1144	11.8249

cobalt(III) chloride resulted in suitable crystals (Fig. 2) that grew to dimensions of up to 400 × 400 × 400 µm and diffracted to 2.5 Å resolution. Accordingly, the final crystallization condition for SBT3 was 0.1 M Bicine pH 8.4, 10% (w/v) PEG 6000, 2.0 M NaCl and 0.01 M hexamminecobalt(III) chloride.

2.4. Data collection and phasing

Prior to data collection, crystals were cryocooled by direct transfer into liquid nitrogen. For cryoprotection of the native crystals, the NaCl concentration was raised to 3.0 M. In the case of the derivative crystals, a fast-soaking procedure was employed. Native crystals were soaked for 20 min or 24 h in the aforementioned cryoprotective solution supplemented with 0.2 M CsCl or 0.01 M TAMM, respectively, followed by direct transfer into liquid nitrogen.

Data sets were collected on Swiss Light Source (SLS) beamline PXII using a MAR225 CCD detector [native: crystal-to-detector distance 230 mm, oscillation range 0.5° in φ , 281 images collected; CsCl derivative: crystal-to-detector distance 150 mm, oscillation range 0.5° in φ , 360 images collected (Fig. 3); TAMM derivative: crystal-to-detector distance 270 mm, oscillation range 0.5° in φ , 120 images collected]. The data were processed and scaled using *XDS* (Kabsch, 1993).

The data sets were combined and scaled using the *CCP4* suite (Collaborative Computational Project, Number, 4, 1994) for input into *SOLVE* (Terwilliger & Berendzen, 1999). Three Cs sites and one TAMM site with an overall Z score of 27.22 and a figure of merit of 0.29 were identified by MIRAS procedures using the INANO and ANOREFINE keywords for the TAMM derivative and INANO and NOANOREFINE for the Cs derivative. *RESOLVE* (Terwilliger, 2000) was then used to obtain an initial electron-density map. Improvement of the electron-density maps and tracing of the protein are now in progress.

The authors are grateful to the staff of the Swiss Light Source (SLS), Villigen, Switzerland for technical support and to Ilme Schlichting, Wulf Blankenfeldt and Eckhard Hofmann for data collection. The work was supported by DFG grant SCHA 591/2 to AS.

References

Berger, D. & Altmann, T. (2000). *Genes Dev.* **14**, 1119–1131.

- Collaborative Computational Project, Number 4 (1994). *Acta Cryst.* **D50**, 760–763.
- Felix, G. & Boller, T. (1995). *Plant J.* **7**, 381–389.
- Fuller, R. S., Brake, A. & Thorner, J. (1989). *Proc. Natl Acad. Sci. USA*, **86**, 1434–1438.
- Kabsch, W. (1993). *J. Appl. Cryst.* **26**, 795–800.
- Liu, J. X., Srivastava, R., Che, P. & Howell, S. H. (2007). *Plant J.* **51**, 897–909.
- Meichtry, J., Amrhein, N. & Schaller, A. (1999). *Plant Mol. Biol.* **39**, 749–760.
- Rautengarten, C., Steinhauser, D., Bussis, D., Stintzi, A., Schaller, A., Kopka, J. & Altmann, T. (2005). *PLoS Comput. Biol.* **1**, e40.
- Siezen, R. J. & Leunissen, J. A. M. (1997). *Protein Sci.* **6**, 501–523.
- Srivastava, R., Liu, J. X. & Howell, S. H. (2008). *Plant J.* **52**, 219–227.
- Terwilliger, T. C. (2000). *Acta Cryst.* **D56**, 965–972.
- Terwilliger, T. C. & Berendzen, J. (1999). *Acta Cryst.* **D55**, 849–861.
- Tripathi, L. & Sowdhamini, R. (2006). *BMC Genomics*, **7**, 200.
- Vera, P. & Conejero, V. (1988). *Plant Physiol.* **87**, 58–63.
- Yamagata, H., Masuzawa, T., Nagaoka, Y., Ohnishi, T. & Iwasaki, T. (1994). *J. Biol. Chem.* **269**, 32725–32731.